

# Mobilization of late-endosomal cholesterol is inhibited by Rab guanine nucleotide dissociation inhibitor

Maarit Hölttä-Vuori\*, Juha Määttä†, Oliver Ullrich‡, Esa Kuismanen† and Elina Ikonen\*

**Cholesterol entering cells in low-density lipoproteins (LDL) via receptor-mediated endocytosis is transported to organelles of the late endocytic pathway for degradation of the lipoprotein particles. The fate of the free cholesterol released remains poorly understood, however. Recent observations suggest that late-endosomal cholesterol sequestration is regulated by the dynamics of lysobisphosphatidic acid (LBPA)-rich membranes [1]. Genetic studies have pinpointed a protein, Niemann-Pick C-1 (NPC-1), that is required for the mobilization of late-endosomal/lysosomal cholesterol by an unknown mechanism [2]. Here, we report the removal of accumulated cholesterol by overexpression of the NPC-1 protein in NPC-1-deficient fibroblasts from patients with Niemann-Pick disease, and in normal fibroblasts upon release of a progesterone-induced block of cholesterol transport. We show that late-endosomal/lysosomal cholesterol mobilization is specifically inhibited by microinjection of Rab GDP-dissociation inhibitor (Rab-GDI). Moreover, clearance of the cholesterol deposits by NPC-1 in patients' fibroblasts is accompanied by the redistribution of LBPA and of a lysosomal hydrolase that utilizes the mannose-6-phosphate receptor. Our results reveal, for the first time, the involvement of a specific molecular component of the membrane-trafficking machinery in cholesterol transport and the coupling of late-endosomal cholesterol egress to the trafficking of other lipid and protein cargo.**

Addresses: \*Department of Biochemistry, National Public Health Institute, Mannerheimintie 166, 00300 Helsinki, Finland. †Department of Biosciences, Division of Biochemistry, Biocenter 1, P.O. Box 56, University of Helsinki, 00014 Helsinki, Finland. ‡Institut für Biochemie, Johannes Gutenberg-Universität, Becherweg 30, D-55218 Mainz, Germany.

Correspondence: Elina Ikonen  
E-mail: elina.ikonen@ktli.fi

Received: 7 September 1999  
Revised: 1 November 1999  
Accepted: 1 December 1999

Published: 14 January 2000

Current Biology 2000, 10:95–98

0960-9822/00/\$ – see front matter  
© 2000 Elsevier Science Ltd. All rights reserved.


## Results and discussion

Transport vehicles carrying LDL-derived cholesterol have not been characterized. Vesicle-mediated transfer would be

consistent with the finding that sterol relocation from lysosomes can be blocked at 21°C [3]. However, several drugs that inhibit vesicular transport, for example cytoskeleton-disrupting agents, were found not to affect the movement of LDL-derived cholesterol (for review see [4]). To address whether the machinery functioning in the membrane trafficking of proteins is involved in cholesterol mobilization, we tested the effect of Rab-GDI, a specific inhibitory protein component of the membrane-trafficking apparatus, on lysosomal/late-endosomal cholesterol egress. Rab-GDI is required for the recycling of Rab GTPases that are involved in vesicle targeting and fusion along the exocytic and endocytic pathways (reviewed in [5]). We prepared recombinant Rab-GDI [6] and verified its activity in removing Rab proteins from membranes. Microinjection of human fibroblasts with Rab-GDI resulted in the redistribution of the late-endosomal marker Rab7 from its punctate perinuclear localization to a diffuse cytosolic localization (Figure 1a,b). Furthermore, Rab-GDI removed up to 75% of Rab proteins from isolated membranes (Figure 1c).


We then established conditions for monitoring cholesterol relocation in LDL-fed cells. Loading of fibroblasts with LDL in the presence of progesterone results in a cholesterol-trafficking block in the late endocytic organelles, which is rapidly reversed upon removal of progesterone [7].

Figure 1



Rab-GDI efficiently removes Rab proteins from membranes. (a,b) Human fibroblasts were microinjected with Rab-GDI and an IgG marker, incubated for 3 h before fixing, and stained with antibodies against (a) IgG and against (b) Rab7. The scale bar represents 20 μm. (c) The ability of Rab-GDI to remove Rab5 from membranes was determined as detailed in the Materials and methods.

Figure 2



Removal of LDL-derived cholesterol is inhibited by microinjection of Rab-GDI in control fibroblasts. (a-f) Cholesterol-starved cells (a) were loaded with LDL in the presence of progesterone (b), and microinjected either with Rab-GDI and IgG (c,d) or with IgG alone (e,f), and further incubated for 3–4 h without progesterone before fixing. The microinjected cells (asterisks) were visualized by fluorescein-conjugated anti-IgG antibodies (c,e) and free cholesterol by filipin staining (a,b,d,f). The scale bar represents 30  $\mu$ m in (a) and 20  $\mu$ m in (c). (g) For quantitation of cholesterol accumulation, non-injected cells (wash-out), cells injected with Rab-GDI and IgG (GDI) and cells injected with IgG (buffer) were scored by filipin staining after the wash-out period. As a control, cells were also stained immediately after LDL and progesterone loading (load). For quantitation of cholesterol accumulation, cells were viewed with a 40 $\times$  objective and the characteristic appearance of multiple bright, perinuclear filipin-positive dots was scored as cholesterol accumulation. The number of cells analyzed from three to five individual experiments, and the standard deviations, are indicated.


We therefore treated fibroblasts with LDL in the presence of progesterone for 24 hours, fixed the cells either directly or after progesterone wash-out, and visualized the accumulation of unesterified cholesterol by light microscopy using the fluorescent antibiotic filipin (Figure 2a,b,d,f). After a wash-out period of 3–4 hours, the discharge of the intense filipin-positive deposits was clearly observed, whereas the faint filipin fluorescence characteristic of untreated cells at the perinuclear area and on the plasma membrane

remained (Figure 2d,f, non-injected cells). When the cells were microinjected with Rab-GDI at the initiation of the wash-out, however, several deposits brightly stained with filipin remained after the wash-out period, indicating a block in cholesterol mobilization (Figure 2c,d). This inhibition was not observed when only the co-injection marker IgG was injected in the same buffer (Figure 2e,f). Quantitation revealed that Rab-GDI inhibited cholesterol relocation by 50% (Figure 2g). Although prolonging the wash-out reduced the number of cells with cholesterol accumulation, the wash-out time was limited by the notion that Rab-GDI loses its effect in the cellular environment with time (data not shown). The degree of transport inhibition by Rab-GDI was in the range previously observed on introduction of GDI into living cells [8–11].

Massive lysosomal deposits of unesterified cholesterol can be found in the fibroblasts of patients suffering from the genetic disease Niemann–Pick C (NPC). These accumulations can be removed by introducing the normal NPC-1 gene product into the cells [2]. We next determined whether the cholesterol transport that resumes in NPC cells after the release of the genetic block is also sensitive to Rab-GDI. NPC-1-deficient fibroblasts were co-transfected with cDNAs for NPC-1 and green fluorescent protein (GFP), and progesterone was added 24 hours after transfection in order to prevent asynchronous cholesterol mobilization from cells expressing slowly increasing levels of NPC-1. Three days after transfection, the transfected cells (identified by GFP fluorescence) were microinjected with Rab-GDI, and cholesterol mobilization was analyzed as above. We found that introduction of Rab-GDI into the transfected cells significantly inhibited complementation of the defect by NPC-1. The cholesterol accumulations remained in 50% of the transfected cells injected with Rab-GDI but in only 20–25% of transfected and control-injected or non-injected cells; the efficiency of transport inhibition by Rab-GDI was comparable to that in normal cells loaded with LDL and progesterone (data not shown). Together, these results suggest that cholesterol is mobilized from the late-endosomal membranes to carriers that rely on Rab proteins for their delivery.

In our experiments, the outcome of removing Rab proteins from their functional cycle by excess Rab-GDI was stagnation of the sterol cargo to the donor membranes. Considering our current understanding of Rab proteins as regulators of the docking and fusion steps of vesicular transport, the anticipated effect would rather have been unsuccessful fusion of transport intermediates to the sites of cargo delivery. The observed retention of the late-endosomal cholesterol might reflect an indirect effect of exhausting the pool of specific Rab proteins required for the transfer process. Moreover, the role of Rab proteins might be more complex than simply involvement in vesicle targeting and fusion. A complex containing Rab5

Figure 3




Redistribution of LBPA upon cholesterol removal in NPC fibroblasts. Confocal micrographs of NPC-1-deficient cells transfected with NPC-1 cDNA and stained as follows. Staining with antibodies against (a) NPC-1 and (b) LBPA; (c) filipin staining of the cells in (a,b). Staining with antibodies against (d) NPC-1 and (e) Lamp-1; (f) filipin staining of cells in (d,e). The scale bar in (a) represents 20  $\mu$ m.

and GDI was recently shown to be required in the vesicle formation stage [12]. Furthermore, GDI inhibits not only vesicular transport but also direct membrane fusion between late endosomes and lysosomes [13].

Immunoelectron microscopy has revealed that some constituents of the late endosomes are confined to the encircling membrane (for example, Lamp-1), whereas others (for example, the mannose-6-phosphate receptor (MPR) and some lysosomal hydrolases that utilize this receptor) localize within the internal membranes [14]. LBPA is also found in the internal membranes [15]. As these membranes are also heavily loaded with cholesterol [16], unpublished observations), we determined whether any other constituents of the late-endosomal membranes would be mobilized along with cholesterol by analyzing the distribution of late-endosomal/lysosomal markers in NPC fibroblasts complemented with the NPC-1 protein. We found that in cells in which the cholesterol accumulation was efficiently removed, the punctate late-endosomal staining typical of LBPA in control and NPC fibroblasts [1] changed dramatically to a diffuse staining pattern (Figure 3a–c). In contrast, the distribution of Lamp-1 was not altered and it continued to co-localize with the NPC-1 protein, which is localized in the late-endosomes/lysosomes handling the sterol load [3]

Figure 4



The redistribution of the lysosomal hydrolase AGA in NPC fibroblasts upon cholesterol removal. (a–f) Confocal micrographs of NPC-1-deficient cells transfected with Myc-tagged NPC-1 cDNA and stained with antibodies against (a) the Myc tag, (c,e,f) against AGA (green) and (d–f) against EEA1 (red). (b) Cholesterol was visualized by filipin staining. (e) Overlay of (c) and (d). (f) The area indicated in (e) at higher magnification. Yellow indicates co-localization (arrows); compare with red EEA1 staining and green AGA staining (arrowheads). The scale bar represents 20  $\mu$ m in (a) and 5  $\mu$ m in (f).

(Figure 3d–f). We also studied the distribution of the soluble lysosomal hydrolase aspartylglucosaminidase (AGA), which uses the mannose 6-phosphate recognition tag [17], in the complemented NPC cells. Interestingly, whereas AGA overlaps extensively with Lamp-1 in both normal fibroblasts and uncorrected NPC cells (data not shown), it became partially redistributed in the corrected NPC cells, as reflected by a change from the characteristic punctate, often perinuclear, ring-like staining to a more scattered pattern (Figure 4a–c). Several of the AGA-positive signals distributed towards the cell periphery co-localized with the early endosomal autoantigen EEA1 (Figure 4d–f), indicating that AGA was partially targeted to early endosomes upon cholesterol mobilization.

No relocation of LBPA and AGA was observed when NPC-1 was overexpressed in normal fibroblasts (data not shown). This suggests that the redistribution of the late endosomal components observed in NPC fibroblasts could be due to missorting or to an alteration in the balance between forward transport and retrieval of cargo in the cholesterol-loaded organelles. The differential sensitivity of the late-endosomal components to redistribution might reflect the hierarchical read-out of their sorting information: whereas Lamp-1 contains a 'dominant' dileucine

motif for lysosomal targeting [18], LBPA might rely on weaker ionic interactions for correct localization.

Our results reveal that late-endosomal cholesterol egress is coupled to the membrane trafficking of other lipid and protein cargo from that compartment. The question then arises as to the destination of the cholesterol-containing carriers. The bulk of lysosomal cholesterol is thought to be mobilized to the plasma membrane [19,20]. There is also evidence for the transit of lysosomally derived cholesterol via the Golgi complex [16]. An alternative route independent of the plasma membrane is, however, suggested to transport cholesterol from the lysosomes to the endoplasmic reticulum [20]. The identification of Rab proteins as regulators of the cholesterol-trafficking circuits now offers the possibility of manipulating these compartment-specific molecular switches to dissect the routes taken by endocytosed cholesterol.

## Materials and methods

The 93.41 NPC fibroblasts were provided by Peter Pentchev (NIH) and F92-99 control fibroblasts by Harriet von Koskull (Helsinki University Central Hospital). For cholesterol loading, control fibroblasts seeded on coverslips were cultured in medium supplemented with 5% lipoprotein-deficient calf serum for 24–72 h. The cells were then transferred into medium supplemented with 50 µg/ml human LDL and 10 µg/ml progesterone for an additional 22–24 h before microinjection. Recombinant His<sub>6</sub>-tagged Rab-GDI (8 mg/ml) was prepared and its activity measured as described previously [6], using the postnuclear supernatant fraction of CHO cells. Rab-GDI was injected into the cytoplasm at 6.4 mg/ml and mouse monoclonal anti-human IgG at 0.8 mg/ml. NPC fibroblasts were seeded on coverslips 1 day before transfection at 50–80% confluency, and transfected using FUGENE6 transfection reagent (Boehringer-Mannheim). For immunofluorescence microscopy, cells fixed with 4% paraformaldehyde were incubated in 10% fetal bovine serum (FBS) supplemented with 0.05% filipin, followed by primary antibodies diluted in 5% FBS/0.05% filipin, and secondary antibodies diluted in 5% FBS. The coverslips were viewed with a Zeiss Axiophot photomicroscope or with a Leica TCS NT confocal microscope.

## Supplementary material

Supplementary material containing more detailed information on the plasmids and antibodies used, and additional methodological details is available at <http://current.biology.com/supmat/supmatin.htm>.

## Acknowledgements

We thank P. Pentchev for providing the NPC fibroblasts, J. Morris for the NPC-1 cDNA, J. Gruenberg for anti-LBPA antibodies, M. Peltola for anti-AGA antibodies, H. Stenmark for anti-EEA1 antibodies, M. Zerial for anti-Rab7 antibodies, R. Keva for technical assistance and V. Olkkonen for help with confocal microscopy. The work was supported by the Academy of Finland (grants 43184, 43668), the Ara Parseghian Medical Research Foundation and the Sigrid Juselius Foundation.

## References

- Kobayashi T, Beuchat M-H, Lindsay M, Frias S, Palmiter RD, Sakuraba H, Parton RG, Gruenberg J: **Late endosomal membranes rich in lysobisphosphatidic acid regulate cholesterol transport.** *Nature Cell Biol* 1999, 1:113-118.
- Carstea ED, Morris JA, Coleman KG, Loftus SK, Zhang D, Cummings C, et al.: **Niemann-Pick C1 disease gene: homology to mediators of cholesterol homeostasis.** *Science* 1997, 277:228-231.
- Neufeld EB, Wastney M, Patel S, Suresh S, Cooney AM, Dwyer NK, et al.: **The Niemann-Pick C1 protein resides in a vesicular compartment linked to retrograde transport of multiple lysosomal cargo.** *J Biol Chem* 1999, 274:9627-9635.
- Liscum L, Munn NJ: **Intracellular cholesterol transport.** *Biochim Biophys Acta* 1999, 1438:19-37.
- Pfeffer SR, Dirac-Svestrup AB, Soldati T: **Rab GDP dissociation inhibitor: putting Rab GTPases in the right place.** *J Biol Chem* 1995, 270:17057-17059.
- Ullrich O, Horiuchi H, Alexandrov K, Zerial M: **Use of Rab-GDP dissociation inhibitor for solubilization and delivery of Rab proteins to biological membranes in streptolysin O-permeabilized cells.** *Methods Enzymol* 1995, 257:243-253.
- Butler JD, Blanchette-Mackie J, Goldin E, O'Neill RR, Carstea G, Roff CF, et al.: **Progesterone blocks cholesterol translocation from lysosomes.** *J Biol Chem* 1992, 267:23797-23805.
- Peter F, Nuoffer C, Pind SN, Balch WE: **Guanine nucleotide dissociation inhibitor is essential for Rab1 function in budding from the endoplasmic reticulum and transport through the Golgi stack.** *J Cell Biol* 1994, 126:1393-1406.
- Ikonen E, Tagaya M, Ullrich O, Montecucco C, Simons K: **Different requirements for NSF, SNAP, and Rab proteins in apical and basolateral transport in MDCK cells.** *Cell* 1995, 81:571-580.
- Funato K, Beron W, Yang CZ, Mukhopadhyay A, Stahl PD: **Reconstitution of phagosome-lysosome fusion in streptolysin O-permeabilized cells.** *J Biol Chem* 1997, 272:16147-16151.
- Chen W, Feng Y, Chen D, Wandinger-Ness A: **Rab11 is required for trans-Golgi network-to-plasma membrane transport and a preferential target for GDP dissociation inhibitor.** *Mol Biol Cell* 1998, 9:3241-3257.
- McLauchlan H, Newell J, Morrice N, Osborne A, West M, Smythe E: **A novel role for Rab5-GDI in ligand sequestration into clathrin-coated pits.** *Curr Biol* 1998, 8:34-45.
- Mullock BM, Bright NA, Fearon CW, Gray SR, Luzio JP: **Fusion of lysosomes with late endosomes produces a hybrid organelle of intermediate density and is NSF dependent.** *J Cell Biol* 1998, 140:591-601.
- Griffiths G, Hoflack B, Simons K, Mellman I, Kornfeld S: **The mannose 6-phosphate receptor and the biogenesis of lysosomes.** *Cell* 1988, 52:329-341.
- Kobayashi T, Stang E, Fang KS, de Moerloose P, Parton RG, Gruenberg J: **A lipid associated with the antiphospholipid syndrome regulates endosome structure and function.** *Nature* 1998, 392:193-197.
- Blanchette-Mackie EJ, Pentchev PG: **Cholesterol distribution in Golgi, lysosomes and endoplasmic reticulum.** In *Intracellular Cholesterol Trafficking*. Edited by Chang TY, Freeman DA. Norwell, MA: Kluwer Academic Publishers; 1998:53-73.
- Ikonen E, Juikonen I, Toilersrud O-K, Kalkkinen N, Peltonen L: **Lysosomal aspartylglucosaminidase is processed to the active subunit complex in the endoplasmic reticulum.** *EMBO J* 1993, 12:295-302.
- Mellman I: **Endocytosis and molecular sorting.** *Annu Rev Cell Dev Biol* 1996, 12:575-625.
- Neufeld EB, Cooney AM, Pitha J, Dawidowicz EA, Dwyer NK, Pentchev PG, Blanchette-Mackie EJ: **Intracellular trafficking of cholesterol monitored with a cyclodextrin.** *J Biol Chem* 1996, 271:21604-21613.
- Underwood KW, Jacobs NL, Howley A, Liscum L: **Evidence for a cholesterol transport pathway from lysosomes to endoplasmic reticulum that is independent of the plasma membrane.** *J Biol Chem* 1998, 273:4266-4274.